

Brain Function Assessment Form™ (BFAF)

Please circle the appropriate number on all questions below. 0 as the least/never to 3 as the most/always.

SECTION 6

- Difficulty with detailed hand coordination 0 1 2 3
- Difficulty with making decisions 0 1 2 3
- Difficulty with suppressing socially inappropriate thoughts 0 1 2 3
- Socially inappropriate behavior 0 1 2 3
- Decisions made based on desires, regardless of the consequences 0 1 2 3
- Difficulty planning and organizing daily events 0 1 2 3
- Difficulty motivating yourself to start and finish tasks 0 1 2 3
- A loss of attention and concentration 0 1 2 3

SECTION 7

- Hypersensitivities to touch or pain 0 1 2 3
- Difficulty with spatial awareness when moving, laying back in a chair, or leaning against a wall 0 1 2 3
- Frequently bumping into the wall or objects 0 1 2 3
- Difficulty with right-left discrimination 0 1 2 3
- Handwriting has become sloppier 0 1 2 3
- Difficulty with basic math calculations 0 1 2 3
- Difficulty finding words for written or verbal communication 0 1 2 3
- Difficulty recognizing symbols, words, or letters 0 1 2 3

SECTION 8

- Difficulty swallowing supplements or large bites of food 0 1 2 3
- Bowel motility and movements slow 0 1 2 3
- Bloating after meals 0 1 2 3
- Dry eyes or dry mouth 0 1 2 3
- A racing heart 0 1 2 3
- A flutter in the chest or an abnormal heart rhythm 0 1 2 3
- Bowel or bladder incontinence, resulting in staining your underwear 0 1 2 3

SECTION 9

- A decrease in movement speed 0 1 2 3
- Difficulty initiating movement 0 1 2 3
- Stiffness in your muscles (not joints) 0 1 2 3
- A stooped posture when walking 0 1 2 3
- Cramping of your hand when writing 0 1 2 3

SECTION 10

- Abnormal body movements (such as twitching legs) 0 1 2 3
- Desires to flinch, clear your throat, or perform some type of movement 0 1 2 3
- Constant nervousness and a restless mind 0 1 2 3
- Compulsive behaviors 0 1 2 3
- Increased tightness and tone in specific muscles 0 1 2 3

SECTION 11

- Difficulty with balance, or balance that is noticeably worse on one side 0 1 2 3
- A need to hold the handrail or watch each step carefully when going down stairs 0 1 2 3
- Episodes of dizziness 0 1 2 3
- Nausea, car sickness, or seasickness 0 1 2 3
- A quick impact after consuming alcohol 0 1 2 3
- A slight hand shake when reaching for something 0 1 2 3
- Back muscles that tire quickly when standing or walking 0 1 2 3
- Chronic neck or back muscle tightness 0 1 2 3